


What claims is:

1. A compound of the general formula I :



I

wherein R¹ is H; C₁~C₄ branched or straight chain alkyl; C₁~C₄ halogenated
 5 branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₄ alkynyl; pyridyl,
 pyrimidinyl, imidazolyl; except H, the above substituents may be
 optionally substituted with one or more following groups: halogen, cyano,
 nitro, hydroxyl, carboxyl, guanidino, C₁~C₄ alkyl, C₁~C₄ alkoxy, C₁~C₄
 alkanoyl, C₃~C₅ cycloalkyl, substituted phenyl, substituted
 10 heterocyclic group, CONR⁵R⁶, NR⁵R⁶, CO₂R⁷, NHSO₂R⁸ or SO₂NR⁹R¹⁰;

R² is H; C₁~C₃ branched or straight chain alkyl; C₁~C₃ halogenated branched
 or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₄ alkynyl; substituted phenyl;
 except H, the above substituents may be optionally substituted with one
 or more following groups: halogen, cyano-, nitro, hydroxyl, carboxyl,
 15 guanidino-, C₁~C₄ alkyl, C₁~C₄ alkoxy, C₁~C₄ alkanoyl, C₃~C₅ cycloalkyl,
 substituted heterocyclic group, CONR⁶R⁷, NR⁶R⁷, CO₂R⁸, NHSO₂R⁹ or
 SO₂NR¹⁰R¹¹;

R³ is H; C₁~C₆ branched or straight chain alkyl which may be optionally
 substituted with C₃~C₆ cycloalkyl or C₁~C₄ alkoxy; C₂~C₄ alkenyl; C₂~C₄
 20 alkynyl;

R⁴ is H; C₁~C₆ branched or straight chain alkyl which may be optionally
 substituted with C₃~C₆ cycloalkyl or C₁~C₄ alkoxy; C₂~C₄ alkenyl; C₂~C₄
 alkynyl;

R⁵ is H; C₁~C₄ branched or straight chain alkyl which may be optionally

substituted with OH, NR^6R^7 , CN, CONR^6R^7 or CO_2R^8 ; $\text{C}_2\sim\text{C}_4$ alkenyl which may be optionally substituted with CN, CONR^6R^7 or CO_2R^8 ; $\text{C}_2\sim\text{C}_4$ alkoxy optionally substituted with NR^6R^7 ; ($\text{C}_2\sim\text{C}_3$ alkoxy) $\text{C}_1\sim\text{C}_2$ branched or straight chain alkyl optionally substituted with OH or NR^6R^7 ; CONR^6R^7 ; 5 CO_2R^8 ; halogen; NR^6R^7 ; $\text{NHSO}_2\text{NR}^6\text{R}^7$; NHSO_2R^9 ; $\text{SO}_2\text{NR}^{10}\text{R}^{11}$; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl, or triazolyl, either of which is optionally substituted with methyl;

R^6 and R^7 are each independently H or $\text{C}_1\sim\text{C}_4$ branched or straight chain alkyl; or R^6 and R^7 together with their attached nitrogen atom form 10 pyrrolinyl, piperidyl, morpholinyl, 4-N(R^{12})-piperazinyl or imidazolyl, either of which is optionally substituted with methyl or hydroxyl;

R^8 is H; $\text{C}_1\sim\text{C}_6$ branched or straight chain alkyl optionally substituted with $\text{C}_1\sim\text{C}_4$ alkoxy, $\text{C}_1\sim\text{C}_4$ alkylamino, dialkylamino; substituted phenyl and substituted heterocyclic group in which the substitut(s) on the ring 15 of substituted phenyl and substituted heterocyclic group are defined as the above;

R^9 is $\text{C}_1\sim\text{C}_3$ alkyl optionally substituted with NR^6R^7 ;

R^{10} and R^{11} are each independently H or $\text{C}_1\sim\text{C}_{12}$ branched or straight chain alkyl; $\text{C}_1\sim\text{C}_3$ halogenated branched or straight chain alkyl; $\text{C}_2\sim\text{C}_6$ alkenyl; 20 $\text{C}_2\sim\text{C}_6$ alkynyl or $\text{C}_3\sim\text{C}_6$ cycloalkyl; or R^{10} and R^{11} take together to form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4-N(R^{13})-piperazinyl; or R^{10} and R^{11} together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4-N(R^{13})-piperazinyl which are optionally substituted with OH, CN, CO_2R^8 , 25 $\text{C}_1\sim\text{C}_4$ branched or straight chain alkyl, $\text{C}_1\sim\text{C}_3$ alkoxy, $\text{NR}^{14}\text{R}^{15}$ or $\text{CONR}^{14}\text{R}^{15}$; substituted phenyl, substituted heterocyclic group, or $\text{C}_1\sim\text{C}_6$ branched or straight chain alkyl substituted with substituted phenyl or substituted heterocyclic group, the said groups are optionally further substituted with OH, CO_2R^8 , $\text{NR}^{14}\text{R}^{15}$, $\text{CONR}^{14}\text{R}^{15}$, or linked together with

another substituted phenyl or substituted heterocyclic group by a carbonyl group;

R^{12} is H; $C_1\sim C_6$ branched or straight chain alkyl which may be optionally substituted with phenyl, $C_2\sim C_3$ alkyl substituted by hydroxyl, or $C_1\sim C_4$ alkoxy; $C_1\sim C_3$ fluoroalkyl; $C_2\sim C_6$ alkenyl; $C_2\sim C_6$ alkynyl; or $C_3\sim C_6$ cycloalkyl; R^{13} is H; $C_1\sim C_6$ branched or straight chain alkyl; $C_2\sim C_6$ branched or straight chain alkyl substituted with $C_1\sim C_3$ alkoxy; $C_2\sim C_6$ branched or straight chain alkyl substituted with hydroxyl; $C_2\sim C_6$ branched or straight chain alkyl substituted with $NR^{14}R^{15}$; $C_2\sim C_6$ branched or straight chain alkyl substituted with phenyl; $C_1\sim C_6$ branched or straight chain alkyl substituted with $CONR^{14}R^{15}$; $C_2\sim C_6$ branched or straight chain hydrocarbyl substituted with CO_2R^8 ; $C_2\sim C_6$ branched or straight chain hydrocarbyl having substituted phenyl or substituted heterocyclic group as substituent; CO_2R^8 , $CONR^{14}R^{15}$, $CSNR^{14}R^{15}$ or $C(NH)NR^{14}R^{15}$; $C_1\sim C_3$ halogenated branched or straight chain alkyl; $C_2\sim C_6$ alkenyl; $C_2\sim C_6$ alkynyl or $C_3\sim C_6$ cycloalkyl; or polyethylene glycol group ($n=2\sim 20$), which is optionally substituted with $C_1\sim C_6$ alkyl on its terminal;

R^{14} and R^{15} are each independently H; $C_1\sim C_4$ branched or straight chain alkyl; $C_2\sim C_4$ branched or straight chain alkyl substituted with $C_1\sim C_3$ alkoxy; or $C_2\sim C_4$ branched or straight chain alkyl substituted with hydroxyl; or R^{14} and R^{15} together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl or morpholinyl; and

the substituted phenyl refers to a phenyl which is substituted with one or more groups selected from $C_1\sim C_4$ alkoxy, halogen, cyano-, CF_3 , OCF_3 , $C_1\sim C_4$ branched or straight chain alkyl on the phenyl ring; The substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms, and the oxides thereof; pentatomic rings containing two or three hetero-atom selected a group consisted of nitrogen, oxygen, and sulfur atoms; the substituting groups on the heterocyclic ring are

C₁~C₄ branched or straight chain alkyl, C₁~C₄ alkoxy, amino, as well as C₁~C₄ branched or straight chain alkyl amino, C₁~C₄ alkoxyamino group; Or their pharmaceutically acceptable salts .

- 5 2.The compound according to claim 1, wherein: R¹ is C₁~C₃ branched or straight chain alkyl optionally substituted with one or more groups selected from a group consisted of the following: C₁~C₄ alkyl, C₁~C₄ alkoxy, C₁~C₄ alkanoyl, substituted phenyl, substituted heterocyclic group, CONR⁶R⁷ and NR⁶R⁷;
- 10 R² is H; C₁~C₃ branched or straight chain alkyl optionally substituted with one or more groups selected from a group consisted of the following: substituted phenyl, substituted heterocyclic group, CONR⁶R⁷, and NR⁶R⁷;
- R³ is H; C₂~C₄ branched or straight chain alkyl which may be optionally substituted with C₃~C₄ cycloalkyl, C₁~C₃ alkoxy; C₂~C₄ alkenyl; or C₂~C₄ alkynyl;
- 15 R⁴ is H; C₁~C₄ branched or straight chain alkyl which may be optionally substituted with C₃~C₅ cycloalkyl or C₁~C₃ alkoxy; C₂~C₄ alkenyl; or C₂~C₄ alkynyl;
- R⁵ is H; C₁~C₄ branched or straight chain alkyl which may be optionally substituted with OH, NR⁶R⁷, CN, CONR⁶R⁷ or CO₂R⁸; C₂~C₄ alkoxy optionally substituted with NR⁶R⁷; NR⁶R⁷; NHSO₂NR⁶R⁷; NHSO₂R⁹; SO₂NR¹⁰R¹¹; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, either of which is optionally substituted with methyl;
- 20 R⁶ and R⁷ are each independently H; C₁~C₄ branched or straight chain alkyl, or R⁶ and R⁷ together with their attached nitrogen atom form a pyrrolinyl, piperidyl, morpholinyl, 4-N(R¹²)-piperazinyl or imidazolyl, either of which is optionally substituted with methyl and hydroxyl;
- 25

R⁸ is H or C₁~C₄ branched or straight chain alkyl;

R⁹ is C₁~C₃ alkyl optionally substituted with NR⁶R⁷;

R¹⁰ and R¹¹ are each independently H or C₁~C₁₂ branched or straight chain alkyl; C₁~C₃ halogenated branched or straight chain alkyl; C₂~C₆ alkenyl;

5 C₂~C₆ alkynyl or C₃~C₆ cylcoalkyl; or R¹⁰ and R¹¹ taken together to form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4-N(R¹³)-piperazinyl; or R¹⁰ and R¹¹ together with their attached nitrogen atom form a pyrrolinyl, pyrrolidone group, piperidyl, morpholinyl, 4-N(R¹³)-piperazinyl; the said groups are optionally substituted with
10 OH, CN, CO₂R⁸, C₁~C₄ branched or straight chain alkyl, C₁~C₃ alkoxyl, NR¹⁴R¹⁵, or CONR¹⁴R¹⁵; substituted phenyl, substituted heterocyclic group, or C₁~C₆ branched or straight alkyl substituted with substituted phenyl or substituted heterocyclic group, the said groups are further substituted with OH, CO₂R⁸, NR¹⁴R¹⁵, CONR¹⁴R¹⁵, or linked together with another
15 substituted phenyl or substituted heterocyclic group by a carbonyl group;

R¹² is H; C₁~C₆ branched or straight chain alkyl which may be optionally substituted with C₂~C₃ alkyl or C₁~C₄ alkoxyl, the said alkyl and alkoxyl are substituted with phenyl, hydroxyl; C₂~C₆ alkenyl or C₃~C₆ cylcoalkyl;

20 R¹³ is H; C₁~C₆ branched or straight chain alkyl; C₂~C₆ branched or straight chain alkyl substituted with C₁~C₃ alkoxyl; C₂~C₆ branched or straight chain alkyl substituted with hydroxyl; C₂~C₆ branched or straight chain alkyl substituted with NR¹⁴R¹⁵; C₂~C₃ branched or straight chain alkyl substituted with phenyl; C₁~C₆ branched or straight chain alkyl
25 substituted with CONR¹⁴R¹⁵; CO₂R⁸, CONR¹⁴R¹⁵, CSNR¹⁴R¹⁵ or C(NH)NR¹⁴R¹⁵; C₁~C₃ halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₆ alkynyl or C₃~C₆ cylcoalkyl;

R¹⁴ and R¹⁵ are each independently H; C₁~C₄ branched or straight chain alkyl; C₂~C₄ branched or straight chain alkyl substituted with C₁~C₃

alkoxyl; or C₂~C₄ branched or straight chain alkyl substituted with hydroxyl; or R¹⁴ and R¹⁵ together with their attached nitrogen atom form pyrrolinyl, pyrrolinone group, piperidyl, or morpholinyl;

The substituted phenyl refers to a phenyl group which is substituted
5 with one or more groups selected from C₁~C₄ alkoxyl, halogen, CN, CF₃, OCF₃, or C₁~C₄ branched or straight chain alkyl; the substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms, and the oxide thereof; or pentatomic rings containing two or three hetero-atom selected a group consisted of nitrogen, oxygen
10 and sulfur atoms; the substituents on the heterocyclic ring are C₁~C₄ branched or straight chain alkyl, C₁~C₄ alkoxyl, amino, as well as C₁~C₄ branched or straight chain alkyl amino, C₁~C₄ alkoxylamino.

3.The compound according to claim 1 or 2, wherein:

R¹ is C₂~C₃ branched or straight chain alkyl which may be optionally
15 substituted with one or more groups selected from substituted heterocyclic group and NR⁶R⁷;

R² is H;

R³ is H; C₂~C₄ branched or straight chain alkyl which may be optionally substituted with C₃~C₄ cycloalkyl; C₂~C₄ alkenyl; C₂~C₄ alkynyl;

20 R⁴ is C₂~C₄ branched or straight chain alkyl, which may be optionally substituted with C₁~C₃ alkoxyl; C₂~C₄ alkenyl; C₂~C₄ alkynyl;

R⁵ is SO₂NR¹⁰R¹¹;

R⁶ and R⁷ together with their attached nitrogen atom form a pyrrolinyl, piperidyl or morpholinyl;

25 R⁸ is H or C₁~C₄ branched or straight chain alkyl;

R¹⁰ and R¹¹ are each independently H or C₁~C₁₂ branched or straight chain alkyl; C₃~C₆ cycloalkyl; or R¹⁰ and R¹¹ take together to form a pyrrolinyl,

pyrrolinone group, piperidyl, morpholinyl, 4-N(R¹³)-piperazinyl; or R¹⁰ and R¹¹ together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, or 4-N(R¹³)-piperazinyl; the said groups are optionally substituted with OH, C₁~C₄ branched or straight chain alkyl, C₁~C₃ alkoxy, NR¹⁴R¹⁵, or CONR¹⁴R¹⁵; substituted phenyl, substituted heterocyclic group, or C₁~C₆ branched or straight alkyl optionally substituted with substituted phenyl, substituted heterocyclic group, the said groups are further substituted with OH, CO₂R⁸, NR¹⁴R¹⁵ or CONR¹³R¹⁴, or linked together with another substituted phenyl or substituted heterocyclic group by a carbonyl;

R¹³ is H; C₁~C₃ branched or straight chain alkyl; C₂~C₃ branched or straight chain alkyl substituted with C₁~C₃ alkoxy; C₂~C₃ branched or straight chain alkyl substituted with OH; C₂~C₆ branched or straight chain alkyl substituted with NR¹⁴R¹⁵; C₂~C₃ branched or straight chain alkyl substituted with phenyl; C₁~C₆ branched or straight chain alkyl substituted with CONR¹⁴R¹⁵; CO₂R⁸ or CONR¹⁴R¹⁵;

R¹⁴ and R¹⁵ are each independently H; C₁~C₄ branched or straight chain alkyl; C₂~C₄ branched or straight chain alkyl substituted with C₁~C₃ alkoxy; or C₂~C₄ branched or straight chain alkyl substituted with OH; or R¹⁴ and R¹⁵ together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl or morpholinyl;

the substituted phenyl refers to a phenyl group which is substituted with one or more substituents selected from a group consisted of C₁~C₄ alkoxy, halogen, CN, CF₃, OCF₃, and C₁~C₄ branched or straight chain alkyl; the substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms and the oxide thereof; or pentatomic rings containing two or three hetero-atom selected a group consisted of nitrogen, oxygen, and sulfur atoms; the substituents on the heterocyclic ring are C₁~C₄ branched or straight chain alkyl, C₁~C₄

alkoxyl, amino, as well as C₁~C₄ branched or straight chain alkyl amino, C₁~C₄ alkoxylamino.

4. The compound according to any one of claims 1-3, wherein the compound is selected from a group consisting of:

5 2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-methoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, and the hydrochloride thereof;

2-[2-n-propoxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, and the hydrochloride thereof;

15 2-[2-allyloxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, and the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-n-propoxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the

monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-ethoxycarbonylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
5 monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-(2-hydroxyethyl)piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride
10 thereof;

2-[2-ethoxyl-5-(pyrrolidinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

15 2-{2-ethoxyl-5-[3-(2-oxy-pyrrolidin-1-yl)-n-propylamino-N-sulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[2-(pyrrolidin-1-yl)-ethylamino-N-sulfonyl]phenyl}-5
20 -methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(morpholino-4-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride,
25 dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(3-(morpholin-4-yl)-n-propylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible

hydrochloride thereof;

2-[2-ethoxyl-5-(2-(morpholin-4-yl)-ethylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(2,6-dimethylmorpholino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

10 2-[2-ethoxyl-5-(1-benzylpiperidyl-4-aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(2-(piperidin-1-yl)ethylamino-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-benzylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-phenylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride

thereof;

2-[2-ethoxyl-5-(4-benzo[1,3]dioxol-5-yl-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[4-(3-phenyl-n-propan-1-yl)piperidyl-1-sulfonyl]phenyl-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(n-propylamino-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(N,N-di(2-hydroxyethyl)aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-methyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-ethyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-n-butyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride

thereof;

2-{2-ethoxyl-5-(p-ethoxylcarboxylphenylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(o-benzoylphenylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(N2-acethydrazido)-N1-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride and dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(2-dimethylaminoethylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-morpholinomethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;


2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-(pyrimidinyl-2)-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride

thereof; and


2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-allyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof.

5.A process for preparing the compounds of general formula I according to claim 1, comprising:

compound IE is reacted with compound IF:




IE



IF


in an inert solvent in the presence of an organic base as a catalyst as well as acid neutralizer at -20°C to 80°C, wherein R¹, R², R³, R⁴ and R⁵ are defined as claim 1, to obtain compound ID:



ID

wherein R¹, R², R³, R⁴ and R⁵ are defined as the above,


then compound IA may be prepared by heating compound ID in an acidic aqueous solution:



IA

wherein R^1 , R^2 , R^3 , R^4 and R^5 are defined as the above,

and then a cyclization reaction of compound IA is carried out by heating
5 and refluxing in an appropriate solution under acidic, basic or neutral
condition to obtain compound of formula I:



I

wherein R^1 , R^2 , R^3 , R^4 and R^5 are defined as the above;

10 or,

compound of formula I is obtained directly by cyclization of the
corresponding compound ID under conditions as follows:


- a) heating to 100~300°C in mixture of phosphorus pentoxide, water and
tertiary amine, or,
- 15 b) reacting at room temperature or reacting by heating in basic hydrogen
peroxide aqueous solution, or,
- c) reacting at room temperature or reacting by heating under acidic
hydrous or anhydrous condition;

or,

specific example of compounds of general formula I wherein R^5 is $SO_2NR^{11}R^{12}$ may be prepared by processes as follows:

compound IC is obtained by reaction between compound IF and compound


5 IE wherein R^5 is H;



IC

wherein R^1 , R^2 , R^3 and R^4 are defined as the above,

then compound IB is obtained by reacting compound IC with chlorosulfonic acid alone or in dichloromethane, chloroform, and other inert or polar
10 non-proton solvents ,




IB

wherein R^1 , R^2 , R^3 and R^4 are as previously defined,

15 and then acylation reaction occurs between compound IB and proper amines in dichloromethane, chloroform, tertiary amine or other inert or polar non-proton solvents at -78°C to 100°C to obtain compound of formula I wherein R^5 is $SO_2NR^{11}R^{12}$, wherein R^{11} and R^{12} are defined as claim 1;

or,

compound IG



wherein R¹, R², R³, R⁴ and R⁵ are defined as claim 1, reacts with compound
 5 IH in the solvents of polar non-proton solvents in the presence of the
 base as catalyst to obtain compound of formula I, wherein X represents
 Cl, Br or I, R³ is defined as claim 1;

Optionally, compounds of the formula I can be converted into the
 corresponding salts by reacting with pharmaceutically acceptable acids.

10 6.A pharmaceutical composition containing the compound according to any
 one of claims 1-7 as active ingredient, and pharmaceutically acceptable
 excipient.


7.A veterinary drugs composition containing the compound according to
 any one of claims 1-7 as active ingredient, and veterinarily acceptable
 15 excipient.

8.Use of compounds according to any one of claims 1-7 for the manufacture
 of a medicament for treatment or prevention of the diseases related with
 phospholipase and the function thereof.


9.The use according to claim 11, wherein the diseases includes: male
 20 sexual (erectile) dysfunction, female sexual dysfunction, premature
 delivery, dysmenorrhea, benign prostatic hyperlasia, bladder
 obstruction, incontinence, stable or unstable angina, hypertension,
 pulmonary hypertension, congestive heart failure, arteriosclerosis,
 stroke, peripheral circulatory disease, low vascular patency, chronic

asthma, allergic asthma, bronchitis, allergic rhinitis, glaucoma, disorder of the gastrointestinal movement, forerunner of the seizure, Kawasaki disease, tolerance of nitric acid ester, multiple sclerosis, peripheral nerve syndrome caused by diabetes, Alzheimer disease (AD), acute respiratory system failure, psoriasis, skin gangrene, metastasis of cancer cell, loss of hair, nutcracker esophagus, anal fissure, and hypoxia-induced vasoconstriction.


10. Intermediates IA-IG for the manufacture of compound of formula I according to claim 1:




IA




IB




IC




ID



IE



IF



IG

2-substituted phenyl-5,7-dihydrocarbyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one derivatives, the preparation and the pharmaceutical use thereof

Field of invention

5 The present invention relates to 2-substituted phenyl-5,7-dihydrocarbyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one derivatives, the process for their preparation, the composition containing them, and the use for treatment and/or prevention of sexual dysfunction and other diseases related to phospholipase 5.

10 Background of invention


Sildenafil, disclosed in WO9428902, is first kind of orally-administrated potent inhibitors of phospholipase 5 in treatment of the erectile dysfunction of man. By inhibiting the phospholipase 5 in corpus cavernosum, it can achieve the purpose of relaxing smooth
15 muscle in human corpus cavernosum, improving penile hyperemia so as to result in erection. The effective rates of sildenafil in treating male sexual organs erectile dysfunction amount to 80%.

Also, Pfizer Ltd. has developed a series of 1,6-dihydropyrrol [4,3-d]pyrimidin-7-one derivatives, and broadened their therapeutic area where
20 such indications was thought to be treated by inhibiting phospholipase 5. All of these compounds are disclosed in EP0951098, WO9849116, US6251904, and WO0024745, and the latter two of patents include the compounds whose substituted phenyl on C-5 is replaced by the substituted pyridin-2-yl. On the basis of the structure of Sildenafil, DONG A PHARMA
25 Co. Ltd. of Korea developed a series of mono substituted derivatives in the nitrogen atom of sulfonylamino group, as disclosed in WO0027848 and WO0198304. Presently, as described in WO0216364, in order to further enhance water-solubility, LG Chem. Invest. Ltd. disclosed the

derivatives of 1,6-dihydropyrrolo[4,3-d]-pyrimidin-7-one with polyethylene glycol. In addition, 1,5-dihydropyrrolo[3,4-d]pyrimidin-4-ones and 1,9-dihydropurin-6-ones were developed by Pfizer Ltd. for the treatment of sexual dysfunction(US6100270). WO0160825 disclosed 3,5-dihydropyrrolo[3,2-d]pyrimidin-4-ones are useful for the treatment of impotence. Recently, 3H-imidazo[5,1-f][1,2,4]triazin-4-ones was disclosed by Bayer Co. Ltd. in the patent application DE19881732.

Detailed description of the invention

The object of the present invention is to provide compounds for treatment of sexual dysfunction and other diseases related to phospholipase 5. Thus, according to one aspect, the invention provides novel aryl substituted 3,7-dihydropyrrolo[2,3-d] pyrimidin-4-ones and their pharmaceutically acceptable salts (also named Yonkenafil), the compounds are the structure of general formula (I):



I

wherein R¹ is H; C₁~C₄ branched or straight chain alkyl; C₁~C₄ halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₄ alkynyl; pyridyl, pyrimidinyl, imidazolyl; except H, the above substituents may be optionally substituted with one or more following groups: halogen, cyano, nitro, hydroxyl, carboxyl, guanidino, C₁~C₄ alkyl, C₁~C₄ alkoxy, C₁~C₄ alkanoyl, C₃~C₅ cycloalkyl, substituted phenyl, substituted heterocyclic group, CONR⁵R⁶, NR⁵R⁶, CO₂R⁷, NHSO₂R⁸ or SO₂NR⁹R¹⁰;

R² is H; C₁~C₃ branched or straight chain alkyl; C₁~C₃ halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₄ alkynyl; substituted phenyl; except H, the above substituents may be optionally substituted with one or more following groups: halogen, cyano-, nitro, hydroxyl, carboxyl, guanidino-, C₁~C₄ alkyl, C₁~C₄ alkoxy, C₁~C₄ alkanoyl, C₃~C₅ cycloalkyl, substituted heterocyclic group, CONR⁶R⁷, NR⁶R⁷, CO₂R⁸, NHSO₂R⁹ or SO₂NR¹⁰R¹¹;

R³ is H; C₁~C₆ branched or straight chain alkyl which may be optionally substituted with C₃~C₆ cycloalkyl or C₁~C₄ alkoxy; C₂~C₄ alkenyl; C₂~C₄ alkynyl;

R⁴ is H; C₁~C₆ branched or straight chain alkyl which may be optionally substituted with C₃~C₆ cycloalkyl or C₁~C₄ alkoxy; C₂~C₄ alkenyl; C₂~C₄ alkynyl;

R⁵ is H; C₁~C₄ branched or straight chain alkyl which may be optionally substituted with OH, NR⁶R⁷, CN, CONR⁶R⁷ or CO₂R⁸; C₂~C₄ alkenyl which may be optionally substituted with CN, CONR⁶R⁷ or CO₂R⁸; C₂~C₄ alkoxy optionally substituted with NR⁶R⁷; (C₂~C₃ alkoxy) C₁~C₂ branched or straight chain alkyl optionally substituted with OH or NR⁶R⁷; CONR⁶R⁷; CO₂R⁸; halogen; NR⁶R⁷; NHSO₂NR⁶R⁷; NHSO₂R⁹; SO₂NR¹⁰R¹¹; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl, or triazolyl, either of which is optionally substituted with methyl;

R⁶ and R⁷ are each independently H or C₁~C₄ branched or straight chain alkyl; or R⁶ and R⁷ together with their attached nitrogen atom form pyrrolinyl, piperidyl, morpholinyl, 4-N(R¹²)-piperazinyl or imidazolyl, either of which is optionally substituted with methyl or hydroxyl;

R⁸ is H; C₁~C₆ branched or straight chain alkyl optionally substituted with C₁~C₄ alkoxy, C₁~C₄ alkylamino, dialkylamino; substituted phenyl and substituted heterocyclic group in which the substitut(s) on the ring

of substituted phenyl and substituted heterocyclic group are defined as the above;

R^9 is C_1 - C_3 alkyl optionally substituted with NR^6R^7 ;

R^{10} and R^{11} are each independently H or C_1 - C_{12} branched or straight chain alkyl; C_1 - C_3 halogenated branched or straight chain alkyl; C_2 - C_6 alkenyl; C_2 - C_6 alkynyl or C_3 - C_6 cycloalkyl; or R^{10} and R^{11} take together to form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4- $N(R^{13})$ -piperazinyl; or R^{10} and R^{11} together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, or 4- $N(R^{13})$ -piperazinyl; the said groups are optionally substituted with OH, CN, CO_2R^8 , C_1 - C_4 branched or straight chain alkyl, C_1 - C_3 alkoxy, $NR^{14}R^{15}$ or $CONR^{14}R^{15}$; substituted phenyl, substituted heterocyclic group, or C_1 - C_6 branched or straight chain alkyl substituted with substituted phenyl or substituted heterocyclic group, the said groups are optionally further substituted with OH, CO_2R^8 , $NR^{14}R^{15}$, $CONR^{14}R^{15}$, or linked together with another substituted phenyl or substituted heterocyclic group by a carbonyl group;

R^{12} is H; C_1 - C_6 branched or straight chain alkyl which may be optionally substituted with phenyl, C_2 - C_3 alkyl substituted by hydroxyl, or C_1 - C_4 alkoxy; C_1 - C_3 fluoroalkyl; C_2 - C_6 alkenyl; C_2 - C_6 alkynyl; or C_3 - C_6 cycloalkyl; R^{13} is H; C_1 - C_6 branched or straight chain alkyl; C_2 - C_6 branched or straight chain alkyl substituted with C_1 - C_3 alkoxy; C_2 - C_6 branched or straight chain alkyl substituted with hydroxyl; C_2 - C_6 branched or straight chain alkyl substituted with $NR^{14}R^{15}$; C_2 - C_6 branched or straight chain alkyl substituted with phenyl; C_1 - C_6 branched or straight chain alkyl substituted with $CONR^{14}R^{15}$; C_2 - C_6 branched or straight chain hydrocarbyl substituted with CO_2R^8 ; C_2 - C_6 branched or straight chain hydrocarbyl having substituted phenyl or substituted heterocyclic group as substituent; CO_2R^8 , $CONR^{14}R^{15}$, $CSNR^{14}R^{15}$ or $C(NH)NR^{14}R^{15}$; C_1 - C_3

halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₆ alkynyl or C₃~C₆ cycloalkyl; or polyethylene glycol group (n=2~20), which is optionally substituted with C₁~C₆ alkyl on its terminal;



R¹⁴ and R¹⁵ are each independently H; C₁~C₄ branched or straight chain alkyl; C₂~C₄ branched or straight chain alkyl substituted with C₁~C₃ alkoxy; or C₂~C₄ branched or straight chain alkyl substituted with hydroxyl; or R¹⁴ and R¹⁵ together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl or morpholinyl; and

the substituted phenyl refers to a phenyl which is substituted with one or more groups selected from C₁~C₄ alkoxy, halogen, cyano-, CF₃, OCF₃, C₁~C₄ branched or straight chain alkyl on the phenyl ring; The substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms, and the oxides thereof; pentatomic rings containing two or three hetero-atoms selected from a group consisting of nitrogen, oxygen, and sulfur atoms; the substituting groups on the heterocyclic ring are C₁~C₄ branched or straight chain alkyl, C₁~C₄ alkoxy, amino, as well as C₁~C₄ branched or straight chain alkyl amino, C₁~C₄ alkoxyamino group.

In another aspect, the invention provides processes for preparation of compounds of the formula I and intermediates used in the preparation thereof.

The process for preparation of compounds of the general formula I comprises:


The compounds of formula IE with a compound of formula IF



IE

IF

wherein R^1 , R^2 , R^3 , R^4 and R^5 are defined as previous, are reacted in an inert solvent such as dichloromethane and toluene and the like to produce the compounds of formula ID,




ID

Wherein R^1 , R^2 , R^3 , R^4 and R^5 are defined as previous.

This reaction is carried out in the presence of an organic base such as tertiary amine, pyridine as a catalyst as well as an acid neutralizer at -20°C to 80°C .


And then compounds of formula IA may be obtained by heating compounds of formula ID in an acidic aqueous solution, generally phosphoric acid aqueous solution:



IA

Wherein R^1 , R^2 , R^3 , R^4 and R^5 are as previously defined.

And then the compounds of formula I is obtained by a cyclization reaction of compounds of formula IA :



I

Wherein R^1 , R^2 , R^3 , R^4 and R^5 are defined as previous.

The cyclization reaction is similar to a conventional method known for
 5 synthesis of pyrimidone. The reaction may be carried out by reflux in
 an appropriate solution under acidic, basic or neutral condition.
 Preferably using alkali metal salts of alcohol or amine, or an organic
 base, and ethanol as a solvent. Thus, for example, the cyclization
 reaction is carried out by refluxing in ethanol and in the presence of
 10 potassium tert-butoxide or sodium ethoxide.

Also, compounds of formula I may be obtained directly by cyclization
 reaction of corresponding compounds of the formula ID.


Generally, this reaction is carried out by heating the formula ID in
 a mixture of P_2O_5 , water and tertiary amine, especially
 15 dimethylcyclohexylamine at 100°C to 300°C .

In an alternative procedure, the reaction may be carried out at room
 temperature or by heating in an alkaline hydrogen peroxide water solution,
 for example a mixture of hydrogen peroxide and urea.

The said reaction may also be carried out at room temperature or by heating
 20 under anhydrous or hydrous acidic conditions generally using
 hydrochloric acid.

The specific example of compounds of formula I wherein R^5 is $\text{SO}_2\text{NR}^{11}\text{R}^{12}$
 may be prepared by following process.


Compounds of formula IC can be readily prepared from compounds of the formula IF and the IE wherein R⁵ is H,



IC

5 wherein R¹, R², R³ and R⁴ are defined as previous.

And then compounds of the formula IB are prepared by the reaction of compounds of the formula IC with chlorosulfonic acid:



IB

10 wherein R¹, R², R³ and R⁴ are defined as previous.


The reaction is generally carried out by heating the formula IC in the presence of an excess amount of chlorosulfonic acid. Also, the said reaction may be performed in solvent of dichloromethane, chloroform, and other inert or polar non-proton solvents. In particular when the
 15 reactant has a poor solubility in chlorosulfonic acid, using above-mentioned solvents seem to be more important. The reaction can be carried out at the temperature as high as 100°C without producing any by-products, but generally in ice bath.

Acylation reaction is carried out by reacting the compounds of the formula
 20 IB with proper amines to obtain the compounds of formula I wherein R⁵

is $\text{SO}_2\text{NR}^{11}\text{R}^{12}$, and R^{11} , R^{12} are defined as previous.

The said acylation reaction may be performed in the solvents of dichloromethane, chloroform, tertiary amine and other inert or polar non-proton solvents at -78°C to 100°C using equal or excess amounts of amines. The excess amount of amine is used not only as a reactant, but also a solvent.

Either, compounds of the formula I are prepared by reacting formula IG with compounds of formula IH, compound IG is compound I wherein R^3 is H:



IG

wherein R^1 , R^2 , R^4 and R^5 are defined as previous.



wherein X represents Cl, Br or I; R^3 is defined as previous.

The said reactions are carried out by heating reflux in the solvents of non-polar proton solvents with organic or non-organic base as catalyst, generally alkali carbonate such as potassium carbonate in ketone solvents such as acetone.

Optionally, compounds of the formula I can be converted into the corresponding salts by reacting with pharmaceutically acceptable acids.

The intermediates of formula IE are prepared from corresponding aromatic carboxylic acid that is commercially available and thionyl chloride,

oxalyl chloride by conventional synthesis method of acyl chloride.

If using oxalyl chloride, the reaction should be carried out in the non-proton solvents such as dichloromethane, chloroform, toluene etc.

at the temperature of -10°C to 60°C for 2-10 hours using an equal or excess

5 (no more than 4 fold) amount of oxalyl chloride in the presence of 0.05-1 equivalent of dimethylformamide as a catalyst. The said reaction solutions may be directly used in the preparation of the formula ID or distilled under the reduced pressure to prepare the purified compounds.

If using thionyl chloride, the reaction should be performed in the
10 non-proton solvents such as dichloromethane, chloroform, toluene etc., or the thionyl chloride itself is used as solvent, and better, under the conditions of reflux for 0.5-3 hours. The obtained solutions may be directly used in the preparation of the formula ID, or distilled under the reduced pressure to prepare the purified compounds.

15 Intermediates of the formula IF used may be synthesized readily from malononitrile and correspondingly substituted 2-aminoketones by conventional synthetic procedures in accordance with literatures described below:

Wiley R.H., et al, J Am. Chem. Soc, 1948. 70, 2005; Johnson R.W. et al,
20 J Heterocyclic Chem. 1997, 14, 383; and Wamhoff H. et al, Synthesis 1976, 51.

The reaction may be carried out in water in the presence of alkali metal hydroxide such as potassium hydroxide or sodium hydroxide etc. as the catalyst at the temperature of 4°C to 60°C . The reaction solution is
25 diluted and filtrated, and dried to obtain the product. If the low water-soluble substituted 2-aminoketones is used, the reaction may be carried out either in two phases of both water and organic reagents in the presence of phase transfer catalysts, or in organic solvents in the

presence of nitrogenous organic base such as pyridine, triethylamine as catalyst. The resulting compounds can be purified by recrystallization.

Another aspect of the invention relates to a pharmaceutical composition for treatment of erectile dysfunction in a male animal, including man, comprising a compound of formula I or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable diluent, excipient or carrier.

Another aspect of this invention relates to a process for the preparation of a pharmaceutical composition for treatment or prevention of erectile dysfunction in a male animal, including man, comprising formulating a compound of formula I or a pharmaceutically acceptable salt thereof with pharmaceutically acceptable diluent, excipient or carrier.

Although the compounds of the invention are envisaged primarily for the treatment of erectile dysfunction or male sexual dysfunction, they may also be useful for the treatment of female sexual dysfunction including orgasmic dysfunction related to clitoral disturbances.

Thus, the aspect of this invention provides the use of a compound of formula I for curing or preventing erectile dysfunction in a male animal, including man, and PDE5-related diseases.

The aforementioned PDE5-related diseases include male sexual (erectile) dysfunction, female sexual dysfunction, premature delivery, dysmenorrhea, benign prostatic hyperlasia, bladder obstruction, incontinence, stable or unstable angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral circulatory disease, low vascular patency, chronic asthma, allergic asthma, bronchitis, allergic rhinitis, glaucoma, disorder of the gastrointestinal movement, forerunner of the seizure, Kawasaki

disease, tolerance of nitric acid ester, multiple sclerosis, peripheral nerve syndrome caused by diabetes, Alzheimer disease (AD), acute respiratory system failure, psoriasis, cutaneous gangrene, metastasis of cancer cell, loss of hair, nutcracker oesophagus, anal fissure, and
5 hypoxia-induced vasoconstriction.


The compounds of the invention can exist in tautomeric forms. It is to be understood that all tautomers and other isomers of formula I as well as the mixture thereof fall into the claimed scope of this invention.

The compounds of this invention may contain one or more asymmetric centers
10 and thus can exist as epimers or optical isomers. Furthermore, they are separated into enantiomers by the conventional methods such as dynamic crystallization or chromatography. Besides, they are synthesized from chiral starting materials or reagents by way of asymmetric synthesis. It is to be understood that all epimers or optical as well as the mixture
15 thereof fall into the claimed scope of this invention.

The compounds of this invention may form pharmaceutically acceptable salts with organic or inorganic acid as well as organic or inorganic alkali.

It is to be understood that this invention includes both mixtures and
20 separate individual pharmaceutically acceptable salts formed by reacting the compounds of this invention with organic or inorganic alkali as well as organic or inorganic acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, citric acid, tartaric acid, gluconic acid, lactic acid, maleic acid, fumaric
25 acid, methane-sulfonic acid, hydroxyacetic acid, succinic acid, 4-toluene sulfonic acid, galacturonic acid, glutamic acid, aspartic acid, and the like.

Preferred compounds of this invention refer to the following compounds of formula I as well as the pharmaceutically acceptable salts thereof:



wherein: R^1 is $C_1\sim C_3$ branched or straight chain alkyl optionally substituted with one or more groups selected from a group consisted of the following: $C_1\sim C_4$ alkyl, $C_1\sim C_4$ alkoxy, $C_1\sim C_4$ alkanoyl, substituted
 5 phenyl, substituted heterocyclic group, $CONR^6R^7$ and NR^6R^7 ;

R^2 is H; $C_1\sim C_3$ branched or straight chain alkyl optionally substituted with one or more groups selected from a group consisted of the following: substituted phenyl, substituted heterocyclic group, $CONR^6R^7$, and NR^6R^7 ;

R^3 is H; $C_2\sim C_4$ branched or straight chain alkyl which may be optionally
 10 substituted with $C_3\sim C_4$ cycloalkyl, $C_1\sim C_3$ alkoxy; $C_2\sim C_4$ alkenyl; or $C_2\sim C_4$ alkynyl;

R^4 is H; $C_1\sim C_4$ branched or straight chain alkyl which may be optionally substituted with $C_3\sim C_5$ cycloalkyl or $C_1\sim C_3$ alkoxy; $C_2\sim C_4$ alkenyl; or $C_2\sim C_4$ alkynyl;

15 R^5 is H; $C_1\sim C_4$ branched or straight chain alkyl which may be optionally substituted with OH, NR^6R^7 , CN, $CONR^6R^7$ or CO_2R^8 ; $C_2\sim C_4$ alkoxy optionally substituted with NR^6R^7 ; NR^6R^7 ; $NHSO_2NR^6R^7$; $NHSO_2R^9$; $SO_2NR^{10}R^{11}$; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, either of which is optionally substituted with methyl;

20 R^6 and R^7 are each independently H; $C_1\sim C_4$ branched or straight chain alkyl, or R^6 and R^7 together with their attached nitrogen atom form a pyrrolinyl, piperidyl, morpholinyl, 4- $N(R^{12})$ -piperazinyl or imidazolyl, either of which is optionally substituted with methyl and hydroxyl;

R^8 is H or $C_1\sim C_4$ branched or straight chain alkyl;

R⁹ is C₁~C₃ alkyl optionally substituted with NR⁶R⁷;

R¹⁰ and R¹¹ are each independently H or C₁~C₁₂ branched or straight chain alkyl; C₁~C₃ halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₆ alkynyl or C₃~C₆ cylcoalkyl; or R¹⁰ and R¹¹ take together to form a
5 pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4-N(R¹³)-piperazinyl; or R¹⁰ and R¹¹ together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, or 4-N(R¹³)-piperazinyl; the said groups the said groups are optionally substituted with OH, CN, CO₂R⁸, C₁~C₄ branched or straight chain alkyl,
10 C₁~C₃ alkoxyl, NR¹⁴R¹⁵, or CONR¹⁴R¹⁵; substituted phenyl, substituted heterocyclic group, or C₁~C₆ branched or straight alkyl substituted with substituted phenyl or substituted heterocyclic group, the said groups are further substituted with OH, CO₂R⁸, NR¹⁴R¹⁵, CONR¹⁴R¹⁵, or linked together with another substituted phenyl or substituted heterocyclic
15 group by a carbonyl group;

R¹² is H; C₁~C₆ branched or straight chain alkyl which may be optionally substituted with C₂~C₃ alkyl or C₁~C₄ alkoxyl, the said alkyl and alkoxyl are substituted with phenyl, hydroxyl; C₂~C₆ alkenyl or C₃~C₆ cylcoalkyl;


R¹³ is H; C₁~C₆ branched or straight chain alkyl; C₂~C₆ branched or straight
20 chain alkyl substituted with C₁~C₃ alkoxyl; C₂~C₆ branched or straight chain alkyl substituted with hydroxyl; C₂~C₆ branched or straight chain alkyl substituted with NR¹⁴R¹⁵; C₂~C₃ branched or straight chain alkyl substituted with phenyl; C₁~C₆ branched or straight chain alkyl substituted with CONR¹⁴R¹⁵; CO₂R⁸, CONR¹⁴R¹⁵, CSNR¹⁴R¹⁵ or C(NH)NR¹⁴R¹⁵; C₁~C₃
25 halogenated branched or straight chain alkyl; C₂~C₆ alkenyl; C₂~C₆ alkynyl or C₃~C₆ cylcoalkyl;

R¹⁴ and R¹⁵ are each independently H; C₁~C₄ branched or straight chain alkyl; C₂~C₄ branched or straight chain alkyl substituted with C₁~C₃ alkoxyl; or C₂~C₄ branched or straight chain alkyl substituted with

hydroxyl; or R^{14} and R^{15} together with their attached nitrogen atom form pyrrolinyl, pyrrolinone group, piperidyl, or morpholinyl;

The substituted phenyl refers to a phenyl group which is substituted with one or more groups selected from $C_1\sim C_4$ alkoxy, halogen, CN, CF_3 , OCF_3 , or $C_1\sim C_4$ branched or straight chain alkyl; the substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms, and the oxide thereof; or pentatomic rings containing two or three hetero-atom selected a group consisted of nitrogen, oxygen and sulfur atoms; the substituents on the heterocyclic ring are $C_1\sim C_4$ branched or straight chain alkyl, $C_1\sim C_4$ alkoxy, amino, as well as $C_1\sim C_4$ branched or straight chain alkyl amino, $C_1\sim C_4$ alkoxyamino.

In the more preferred embodiment, compounds of the general formula I and their pharmaceutically acceptable salts as follows:



I

wherein R^1 is $C_2\sim C_3$ branched or straight chain alkyl which may be optionally substituted with one or more groups selected from substituted heterocyclic group and NR^6R^7 ;

R^2 is H;

R^3 is H; $C_2\sim C_4$ branched or straight chain alkyl which may be optionally substituted with $C_3\sim C_4$ cycloalkyl; $C_2\sim C_4$ alkenyl; $C_2\sim C_4$ alkynyl;

R^4 is $C_2\sim C_4$ branched or straight chain alkyl, which may be optionally substituted with $C_1\sim C_3$ alkoxy; $C_2\sim C_4$ alkenyl; $C_2\sim C_4$ alkynyl;

R^5 is $SO_2NR^{10}R^{11}$;

R^6 and R^7 together with their attached nitrogen atom form a pyrrolinyl, piperidyl or morpholinyl;

R^8 is H or $C_1\sim C_4$ branched or straight chain alkyl;

5 R^{10} and R^{11} are each independently H or $C_1\sim C_{12}$ branched or straight chain alkyl; $C_3\sim C_6$ cycloalkyl; or R^{10} and R^{11} take together to form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, 4-N(R^{13})-piperazinyl; or R^{10} and R^{11} together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl, morpholinyl, or 4-N(R^{13})-piperazinyl; the
10 said groups are optionally substituted with OH, $C_1\sim C_4$ branched or straight chain alkyl, $C_1\sim C_3$ alkoxy, $NR^{14}R^{15}$, or $CONR^{14}R^{15}$; substituted phenyl, substituted heterocyclic group, or $C_1\sim C_6$ branched or straight alkyl optionally substituted with substituted phenyl, substituted heterocyclic group, the said groups are further substituted with OH,
15 CO_2R^8 , $NR^{14}R^{15}$ or $CONR^{13}R^{14}$, or linked together with another substituted phenyl or substituted heterocyclic group by a carbonyl;

R^{13} is H; $C_1\sim C_3$ branched or straight chain alkyl; $C_2\sim C_3$ branched or straight chain alkyl substituted with $C_1\sim C_3$ alkoxy; $C_2\sim C_3$ branched or straight chain alkyl substituted with OH; $C_2\sim C_6$ branched or straight chain alkyl
20 substituted with $NR^{14}R^{15}$; $C_2\sim C_3$ branched or straight chain alkyl substituted with phenyl; $C_1\sim C_6$ branched or straight chain alkyl substituted with $CONR^{14}R^{15}$; CO_2R^8 or $CONR^{14}R^{15}$;

R^{14} and R^{15} are each independently H; $C_1\sim C_4$ branched or straight chain alkyl; $C_2\sim C_4$ branched or straight chain alkyl substituted with $C_1\sim C_3$
25 alkoxy; or $C_2\sim C_4$ branched or straight chain alkyl substituted with OH; or R^{14} and R^{15} together with their attached nitrogen atom form a pyrrolinyl, pyrrolinone group, piperidyl or morpholinyl;

the substituted phenyl refers to a phenyl group which is substituted

with one or more substituents selected from a group consisted of C₁~C₄ alkoxyl, halogen, CN, CF₃, OCF₃, and C₁~C₄ branched or straight chain alkyl; the substituted heterocyclic group refers to hexatomic rings containing one or two nitrogen atoms and the oxide thereof; or pentatomic rings containing two or three hetero-atom selected a group consisted of nitrogen, oxygen, and sulfur atoms; the substituents on the heterocyclic ring are C₁~C₄ branched or straight chain alkyl, C₁~C₄ alkoxyl, amino, as well as C₁~C₄ branched or straight chain alkyl amino, C₁~C₄ alkoxyamino.

Especially preferred compounds of invention include:

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-methoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-n-propoxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-allyloxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, and the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-n-propoxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the

monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
5 monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride
10 thereof;

2-[2-ethoxyl-5-(4-ethoxycarbonylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride
thereof;

15 2-[2-ethoxyl-5-(4-(2-hydroxyethyl)piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride
thereof;

2-[2-ethoxyl-5-(pyrrolidinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl
20 -3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride,
dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[3-(2-oxy-pyrrolidin-1-yl)-n-propylamino-N-sulfonyl]
phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-
one, the monohydrochloride, dihydrochloride and other possible
25 hydrochloride thereof;

2-{2-ethoxyl-5-[2-(pyrrolidin-1-yl)-ethylamino-N-sulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
monohydrochloride, dihydrochloride and other possible hydrochloride

thereof;

2-[2-ethoxyl-5-(morpholino-4-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

5 2-[2-ethoxyl-5-(3-(morpholin-4-yl)-n-propylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

10 2-[2-ethoxyl-5-(2-(morpholin-4-yl)-ethylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

15 2-[2-ethoxyl-5-(2,6-dimethylmorpholino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

20 2-[2-ethoxyl-5-(1-benzylpiperidyl-4-aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(2-(piperidin-1-yl)ethylamino-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

25 2-[2-ethoxyl-5-(4-benzylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-phenylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

5 2-[2-ethoxyl-5-(piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-benzo[1,3]dioxol-5-yl-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[4-(3-phenyl-n-propan-1-yl)piperidyl-1-sulfonyl]phenyl-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(n-propylamino-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

20 2-[2-ethoxyl-5-(N,N-di(2-hydroxyethyl)aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-methyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-ethyl]aminosulfonyl}phenyl}-5-

methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-n-butyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(p-ethoxylcarboxylphenylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(o-benzoylphenylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(N2-acethydrazido)-N1-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride and dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(2-dimethylaminoethylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-{2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl}-5-morpholino

methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-(pyrimidinyl-2)-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof;

and

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-allyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the monohydrochloride, dihydrochloride and other possible hydrochloride thereof.

The compounds of general formula I can not only be prepared into orally-administrated solid formulations, such as the tablets, pills, capsules and powder, but also liquid ones, such as suspensions, solution, emulsion and syrup. All of these formulations may comprise a variety of conventional excipients, such as the wetting agents, sweet-enhancers, aromatics and preservatives, etc., they may also comprise some other conventional functional excipients, such as the fillers (starch and carbohydrates), binders (carboxymethylcellulose etc.), dispersants (calcium carbonate and sodium carbonate etc.), diluents (glycerol), absorption enhancers (quaternary ammonium compounds), lubricants (stearate) and absorption agents (kaolin).

The compounds of the formula I can be prepared into ointment for external use. Likewise, they can be also prepared into intravenous injections.

Generally, for human, oral administration of the compounds of this invention is the preferred route, being the most convenient route and avoiding the disadvantages associated with administration in corpus

cavernosum. In circumstances where the patients suffers from a swallowing disorder or from impairment of drug absorption after oral administration, the drug may be administrated parenterally, e.g. sublingually, buccally, transdermally or injection.

5 For veterinary use, a compound of formula I or a non-toxic salt thereof is administered as a suitable acceptable formulation in accordance with common veterinary practice and the veterinary surgeon will determine the dose range and route of administration, which will be the most appropriate for a particular male animal.

10 Furthermore, none of any obvious sign of adverse acute toxicity is shown for the compounds of this invention tested in rat and dog, both intravenously (i.v.) and orally (p.o.) at up to 3 mg/Kg, has shown. For the situation of mice, no deaths occurred after doses of up to 100mg/Kg i.v.. The LD₅₀ for a single dose of compound I-HCl in mice is 2000mg/kg.


15 **Best modes for carrying out the invention**

Now, the preparing methods of the compounds of the present invention are further illustrated by the preparation of 2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, the
20 monohydrochloride and dihydrochloride thereof as example.

Example 1

Preparation of 2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one, its monohydrochloride and dihydrochloride

25 Route of synthesis



(1a) 2-amino-3-cyano-4-methylpyrrole;

5 (1b) N-propyl-2-amino-3-cyano-4-methylpyrrole;

(2) 2-ethoxylbenzoyl chloride;

(3a) N-(3-cyano-4-methyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide;

(3b) N-(3-cyano-4-methyl-1-n-propyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide;

5 (4a) 2-(2-ethoxylbenzamido)-4-methyl-1H-pyrrolo-3-formamide;

(4b) 2-(2-ethoxylbenzamido)-4-methyl-1-n-propyl-1H-pyrrolo-3-formamide;

(5) 2-(2-ethoxylphenyl)-5-methyl-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one;

10 (6) 2-(2-ethoxylphenyl)-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one;

(7) 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonyl chloride;

15 (8) 2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one.

Preparation 1

N-(3-cyano-4-methyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide (3a) and N-(3-cyano-4-methyl-1-n-propyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide (3b):

20 2-ethoxyl benzoic acid (10.0g, 60.2mmol) was added into thionyl chloride (20ml), and the mixture was refluxed with agitation for 40 minutes, and the excess amount of thionyl chloride was evaporated under reduced pressure. The residual was dissolved into dichloromethane (150ml).

25 Within 30 minutes and being stirred on ice bath, the afore-obtained solution of 2-ethoxyl benzoyl chloride was dropped into the compound (1a) (7.0g, 56.8mmol) dissolved in tetrahydrofuran (80ml) and

triethylamine (8.5ml, 61.0mmol). After completion, the mixture was stirred for 1 hour at 0°C. After being washed with water and filtrated with diatomaceous earth, the reaction solution was mixed with 20g of silica gel and evaporated to dryness. The resulting residual was eluted with dichloromethane by using silica gel(80g) column to obtain 7.5g of solid product(3a) with the yield of 48%. Furthermore, the sample for analysis was prepared by column chromatography (developing agent: dichloromethane: n-hexane=1:2) and recrystallization (dichloromethane: n-hexane=1:5).

mp 183~184°C (sublimation 162°C);

IR (cm⁻¹): 3326, 3309, 2981, 2938, 2915, 2854, 2208, 1647, 1593, 1471, 1309, 1302, 1232, 1039, 923, 727, 655, 648;

¹H NMR (CDCl₃): δ 1.70 (t, J=7.0Hz, 3H), 2.15 (s, 3H), 4.32 (q, J=7.0Hz, 2H), 6.24 (s, 1H), 7.04 (d, 1H), 7.10 (m, 1H), 7.51 (dd, 1H), 8.20 (dd, J=7.9 and 1.8Hz, 1H), 10.69 (brs, 1H), 10.80 (s, 1H);

¹³C NMR (CDCl₃): δ (CH₃) 10.6, 15.0; (CH₂) 65.7; (CH) 110.3, 112.3, 121.4, 132.1, 134.2; (C) 78.7, 115.6, 119.2, 119.4, 136.7, 157.0, 163.2;

MS (ES⁺): m/z 287 (M+NH₄).

Elemental analysis (C₁₅H₁₅N₃O₂): C 66.90%; H 5.61%; N 15.60%; O 11.88%.

The compound (3b) was prepared from compound(1b) according to the above-mentioned method with the yield of 41%.

mp 58~61°C;

IR (cm⁻¹): 3596, 3336, 2969, 2937, 2877, 2216, 1676, 1658, 1603, 1571, 1537, 1475, 1431, 1292, 1232, 1122, 1037, 927, 789, 752, 577;

¹H NMR (CDCl₃): δ 0.88 (t, J=7.4Hz, 3H), 1.58 (t, J=7.0Hz, 3H), 1.75 (m,

2H), 2.16 (s, 3H), 3.73 (t, J=7.4Hz, 2H), 4.30 (q, J=7.0Hz, 2H), 6.36 (s, 1H), 7.04 (d, 1H), 7.11 (m, 1H), 7.48 (dd, 1H), 8.23 (dd, J=7.9 and 1.8Hz, 1H), 9.62 (brs, 1H);

^{13}C NMR (CDCl_3): δ (CH₃) 11.1, 14.8; (CH₂) 23.6, 48.3, 65.2; (CH) 112.5, 117.0, 121.3, 132.5, 134.1; (C) 89.2, 115.6, 119.8, 120.5, 131.2, 157.1, 165.0;

MS (ES^+): m/z 329 ($\text{M}+\text{NH}_4$).

Preparation 2

2-(2-ethoxylbenzamido)-4-methyl-1H-pyrrolo-3-formamide (4a) and
2-(2-ethoxylbenzamido)-4-methyl-1-n-propyl-1H-pyrrolo-3-formamide (4b);

A mixture of N-(3-cyano-4-methyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide (3a) (2.00g, 7.44mmol) or N-(3-cyano-4-methyl-1-n-propyl-1H-pyrrol-2-yl)-2-ethoxylbenzamide (3b) (2.30g, 7.44mmol) of preparation 1 and 85% phosphoric acid (14.8ml) was stirred for 20 minutes at 130°C, cooled and poured into crushed ice (80g). The precipitations were filtrated and dried to give dark red solid of compound (3a) or (3b) with the yield of 80%. The product (3a) and (3b) of this step may be directly used for the next step without further purification.

Preparation 3

2-(2-ethoxylphenyl)-5-methyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one (5) and 2-(2-ethoxylphenyl)-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one (6)

A mixture of 2-(2-ethoxylbenzamido)-4-methyl-1H-pyrrolo-3-formamide (4a) (7.0g, 25.5mmol) of preparation 2 and dimethyl cyclohexylamine (20ml) was refluxed with agitation for 11 hours in N,N-dimethyl formamide

(100ml). After evaporation the solvent by distillation under reduced pressure, the residual was extracted with dichloromethane, and the dichloromethane extraction was washed with water. the resultant extraction was dried with anhydrous sodium sulfate. n-hexane (80ml) was added into the residual and ground to give product (5) (6.0g) by filtration with the yield of 91%.

mp 219~221°C

IR (cm⁻¹): 3187, 3114, 3062, 2978, 2923, 1658, 1587, 1460, 1321, 1292, 1250, 1044, 771, 763;

¹H NMR (DMSO-d₆): δ 1.35 (t, J=6.9Hz, 3H), 2.29 (s, 3H), 4.13 (q, J=7.0Hz, 2H), 6.79 (s, 1H), 7.05 (t, 1H), 7.14 (d, 1H), 7.45 (dd, 1H), 7.76 (dd, 1H), 11.35 (brs, 1H), 11.54 (brs, 1H);

¹³C NMR (DMSO-d₆): δ (CH₃) 11.2, 14.5; (CH₂) 64.2; (CH) 113.0, 118.0, 120.6, 130.1, 131.9, (C) 105.0, 113.6, 121.9, 148.5, 149.8, 156.5, 159.2;

MS(ES⁺): m/z 287 (M+NH₄).

The compound (6) was prepared from compound(4b) according to the above-mentioned method with the yield of 85%.

mp 124~127°C

IR (cm⁻¹): 3234, 3184, 3141, 3103, 3056, 2956, 2943, 2869, 1654, 1595, 1567, 1468, 1311, 1267, 1243, 1191, 1118, 1047, 758;

¹H NMR (CDCl₃): δ 0.88 (t, J=7.5Hz, 3H), 1.23 (t, 3H), 1.80 (q, 2H), 2.42 (s, 3H), 4.08 (t, J=7.2Hz, 2H), 4.22 (q, 2H), 6.60 (s, 1H), 7.01 (d, J=8.3Hz, 1H), 7.08 (t, 1H), 7.40 (m, 1H), 8.35 (dd, J=8.0 and 1.9 Hz, 1H), 11.02 (brs, 1H).

Preparation 4

2-(2-ethoxylphenyl)-5-methyl-7-n-propyl-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one(6):

A mixture of compound (5) (1.5g, 5.57mmol) of preparation 3, n-propyl bromide (2.0g, 16.3mmol) and potassium carbonate (5g, 36.2mmol) was dissolved in acetone (15ml), refluxed with agitation by heating for 15 hours, after the solids were filtrated out, the filtrate was dried under reduced pressure. The resultant was developed by column chromatography, using dichloromethane as mobile phase to obtain 0.6g of product(6) with yield of 35%. The physical/chemical data were identical with that of the above-mentioned.

Preparation 5

4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonyl chloride(7):

2-(2-ethoxylphenyl)-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one(6) (1.25g, 4.01mmol) of preparation 4 was added into chlorosulfonic acid (4ml) that was dissolved in acetic ether (20ml), stirred at 0°C by two batches. The obtained solution was stirred at 0 °C for 30 minutes, and then reacted with agitation at room temperature for 3 hours. The resultant solution was poured into the a mixture of icy water (50ml) and acetic ether (50ml). The organic layer was separated, washed with cold water (5ml), desiccated with anhydrous sodium sulfate and concentrated to dryness to afford 1.33g of product as yellow foam. The yield was 81%. The product was used directly for the next reaction.

Compound 1:

2-[2-ethoxyl-5-(4-ethyl-piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one (8):

4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydro-3H-pyrrolo[2,3-d]

]pyrimidin-2-yl)benzenesulfonyl chloride (7) (1.00g, 2.44mmol) of Preparation 5 was dissolved into dichloromethane (20ml), stirred at 0 °C, into which 1-ethyl piperazine (0.78ml, 6.10mmol) was added slowly. Reactant solution was stirred at 0 °C for 5 minutes, and then sequentially stirred at room temperature for 5 hours. The crude product was washed with water and dried with anhydrous sodium sulfate to give 1.2g of product as yellow foam. Continuously, the product was refined by column chromatography (acetic ether: methanol=20:1) to afford 0.89g of product as a yellow solid with yield of 75%.

10 mp: 174~176 °C (EtOAc);

IR (cm⁻¹): 3324, 2960, 2923, 2869, 2862, 2767, 1682, 1560, 1458, 1355, 1282, 1247, 1172, 1149, 739, 615, 588, 555;

¹H NMR (CDCl₃): δ 0.89 (t, J=7.4Hz, 3H), 0.99 (t, J=7.2Hz, 3H), 1.61 (t, J=7.0Hz, 3H), 1.77-1.86 (m, 2H), 2.35 (m, 2H), 2.41 (s, 3H), 2.50 (brs, 4H), 3.05 (brs, 4H), 4.08 (t, J=7.0Hz, 2H), 4.29-4.37 (q, 2H), 6.61 (s, 1H), 7.11 (d, J=8.8Hz, 1H), 7.77 (dd, J=8.7 and 2.2Hz, 1H), 8.74 (d, J=2.2, 1H), 10.63 (brs, 1H);

¹³C NMR (CDCl₃): δ (CH₃) 11.0, 11.3, 11.8, 14.3; (CH₂) 23.8, 45.9, 46.1, 51.6, 51.7, 65.8; (CH) 112.9, 121.1, 130.6, 131.3; (C) 105.7, 114.6, 121.4, 127.8, 146.8, 147.3, 159.3, 159.6;

20 MS (ES⁺): m/z 505 (M+NH₄).

Elemental analysis (C₂₄H₃₃N₅O₄S): theoretical value C 59.12%; H 6.82%; N 14.36%; practically measured value C 59.38%; H 7.10%; N 14.12%.

Compound 1-HCl:

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one monohydrochloride

(9):

The free alkali (compound 1) (1.00g, 2.05mmol) was dissolved into ether (10ml) and dichloromethane (10ml), into which the solution of 4M hydrochloric acid (HCl)- dioxane (0.51ml, 2.04mmol) diluted with ethyl ether (10ml) was dropped with agitation. After completion, the resulting solution was continued to stir at room temperature for 20 minutes, filtrated and dried to give 1.01g of monohydrochloride with yield of 94%.

mp: 147~150°C;

IR(cm^{-1}): 2964, 2931, 2675, 2599, 2462, 1668, 1574, 1456, 1348, 1167, 933, 588;

^1H NMR(D_2O): δ 0.72(m, 3H) ,1.24(t, $J=7.3\text{Hz}$, 3H), 1.45(m, 3H), 1.59(m, 2H), 2.04(s, 3H), 2.77-3.81(all brs, 8H), 3.20(q, 2H), 3.75(m, 2H), 4.20(m, 2H), 6.62(m, 1H), 7.17(m, 1H), 7.73(m, 1H), 8.22(s, 1H).

Elemental analysis ($\text{C}_{24}\text{H}_{33}\text{N}_5\text{O}_4\text{S} \cdot \text{HCl}$): theoretical value C 55.00%; H 6.54%; N 13.36%; practically measured value C55.28%; H 6.41%; N 13.07%.

Compound 1-2HCl:

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one dihydrochloride

mp: 177~180°C;

IR(cm^{-1}): 2962, 2929, 2677, 2597, 2456, 1652, 1569, 1458, 1357, 1276, 1162, 1093, 1027, 939, 731, 582;

^1H NMR(D_2O): δ 0.64(t, $J=7.4\text{Hz}$, 3H), 1.23(t, $J=7.3\text{Hz}$, 3H), 1.40(t, $J=6.9\text{Hz}$, 3H), 1.51(m, 2H), 1.98(s, 3H), 2.74(m, 2H), 3.12(m, 2H), 3.19(t, 2H), 3.56(m, 2H), 3.65(t, 2H), 3.78(d, 2H), 4.12(q, 2H), 6.43(s, 1H), 7.10(d,

J=9.1Hz, 1H) , 7.68(dd, J=8.8 and 2.3Hz, 1H), 8.16(d, J=2.3Hz, 1H).

Likewise, by using similar processes, the starting materials were correspondingly used to synthesize the compounds as follows:

Compound 2:

5 2-[2-methoxyl-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

Compound of
4-methoxyl-3-(5-methyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared by using the
10 2-amino-3-cyano-4-methylpyrrole (IE) and 2-methoxylbenzoyl chloride (IF) as starting materials via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

¹H NMR(CDCl₃): δ 0.86(t, 3H), 0.99(t, 3H), 1.73(m, 2H), 2.32(q, 2H), 2.40(s, 3H), 2.48(brs, 4H), 3.04(brs, 4H), 4.08(t, 2H), 4.14(s, 3H), 6.60(s, 1H),
15 7.0(d, 1H), 7.74(dd, 1H), 8.77(d, 1H).

MS(ES): m/z 491 (M+NH₄).

Compound 3:

2-[2-n-propyloxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-
20 7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-n-propyloxy-3-(5-methyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared by using
2-amino-3-cyano-4-methylpyrrole (IE) and 2-n-propyloxybenzoyl chloride (IF) as starting materials via the corresponding intermediates
25 ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

¹H NMR(CDCl₃) δ: 0.87(t, 3H), 1.00(t, 3H), 1.16(t, 3H), 1.79(m 2H), 2.01(m, 2H), 2.36(q, 2H), 2.42(s, 3H), 2.50(brs, 4H), 3.04(brs, 4H), 4.08(t, 2H), 4.21(t, 2H), 6.60(s, 1H), 7.09(d, 1H), 7.77(dd, 1H), 8.77(d, 1H).

5 MS(ES⁺): m/z 519 (M+NH₄).

Compound 4:

2-[2-allyloxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-allyloxy-3-(5-methyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared by using
10 2-amino-3-cyano-4-methylpyrrole (IE) and 2-allyloxybenzoyl chloride (IF) as starting materials via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

15 ¹H NMR(CDCl₃) δ: 0.83(t, 3H), 0.97(t, 3H), 1.82(t, 2H), 2.37(q, 2H), 2.41(s, 3H), 2.52(brs, 4H), 3.07(brs, 4H), 4.07(t, 2H), 4.62(m, 2H), 5.24(m, 2H), 5.83(m, 1H), 6.62(s, 1H), 7.12(d, 1H), 7.79(dd, 1H), 8.69(d, 1H), 9.97(br, 1H).

MS(ES⁺): m/z 517 (M+NH₄).

20 Compound 5:

2-[2-n-propyloxy-5-(4-ethylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-n-propyloxy-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) was prepared by using
25 2-amino-3-cyano-4-ethylpyrrole (IE) and 2-propyloxybenzoyl chloride (IF) as starting materials via the corresponding intermediates ID, IA,

IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

¹H NMR(CDCl₃) δ : 0.88 (t, 3H), 0.99 (t, 3H), 1.04 (t, 3H), 1.52 (t, 3H), 1.74-1.84 (m, 4H), 2.35 (q, 2H), 2.46 (brs, 4H), 2.68 (q, 2H), 3.05 (brs, 4H), 4.09 (t, 2H), 4.36 (t, 2H), 6.61 (s, 1H), 7.09 (d, 1H), 7.81 (dd, 1H), 8.71 (d, 1H).

MS (ES⁺): m/z 533 (M+NH₄).

Compound 6:

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) obtained from previous Preparation 5 and 1-methylpiperazine.

¹H NMR(D₂O): δ : 0.90 (t, 3H), 1.62 (s, 3H), 1.78 (m, 2H), 2.15 (s, 3H), 2.41 (s, 3H), 2.48 (brs, 4H), 3.04 (brs, 4H), 4.08 (t, 2H), 4.34 (q, 2H), 6.61 (s, 1H), 7.09 (d, 1H), 7.74 (dd, 1H), 8.76 (d, 1H).

MS (ES): m/z 491 (M+NH₄).

Compound 7:

2-[2-ethoxyl-5-(4-methylpiperazinyl-1-sulfonyl)phenyl]-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-ethyl-4-chloro-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) was prepared by using 2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxylbenzoyl chloride (IF) as starting materials via the corresponding intermediates ID, IA, IG

(IH) and IC. The resulting compound was reacted with 1-methylpiperazine to give the titled compound.

¹H NMR(CDCl₃) δ: 0.92 (t, 3H), 1.54-1.64 (m, 6H), 1.75 (m, 2H), 2.14 (s, 3H), 2.53 (m, 6H), 3.04 (brs, 4H), 4.09 (t, 2H), 4.37 (q, 2H) 6.62 (s, 1H), 7.11 (d, 1H), 7.80 (dd, 1H), 8.76 (d, 1H).

MS(ES): m/z 505 (M+NH₄).

Compound 8:

2-[2-ethoxyl-5-(4-ethoxylcarbonylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) obtained from previous Preparation 5 and 1-ethoxylcarbonylpiperazine.

¹H NMR(CDCl₃) δ: 0.90 (t, 3H), 1.25 (t, 3H), 1.62 (t, 3H), 1.83 (m, 2H), 2.41 (s, 3H), 2.48 (brs, 4H), 3.07 (brs, 4H), 4.05 (q, 2H), 4.32 (q, 2H), 4.41 (q, 2H), 6.62 (s, 1H), 7.12 (d, 1H), 7.81 (d, 1H), 8.67 (s, 1H).

MS(ES⁺): m/z 549 (M+NH₄).

Compound 9:

2-[2-ethoxyl-5-(4-(2-hydroxyethyl)piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and 1-(2-hydroxyethyl)piperazine.

¹H NMR(CDCl₃) δ: 0.89 (t, 3H), 1.61 (t, 3H), 1.72 (m, 2H), 2.41 (s, 3H),

2.49-2.58(m, 6H), 3.07(brs, 4H), 3.67(m, 2H), 4.08(t, 2H), 4.33(q, 2H), 6.61(s, 1H), 7.10(d, 1H), 7.76(dd, 1H), 8.69(d, 1H).

MS(ES⁺): m/z 521 (M+NH₄).

Compound 10:

5 2-[2-ethoxyl-5-(pyrrolidinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and
10 pyrrolidine.

¹H NMR(CDCl₃) δ: 0.89(t, 3H), 1.61(t, 3H), 1.81(m, 6H), 2.41(s, 3H), 3.25(m, 4H), 4.10(t, 2H), 4.34(q, 2H), 6.62(s, 1H), 7.12(d, 1H), 7.85(dd, 1H), 8.83(t, 1H), 10.76(brs, 1H).

MS(ES⁺): m/z 462 (M+NH₄).

15 Compound 11:

2-{2-ethoxyl-5-[3-(2-oxy-pyrrolidinyl-1)-n-propylamino-N-sulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of
20 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) of Preparation 5 and 3-(2-oxy-pyrrolidinyl-1)-n-propylamine.

¹H NMR(CDCl₃) δ: 0.90(t, 3H), 1.55(t, 2H), 1.68(m, 2H), 1.78(m, 2H), 1.94(t, 2H), 2.27(t, 2H), 2.38(s, 3H), 2.87(t, 2H), 3.28(m, 4H) 4.07
25 (t, 2H), 4.30(t, 2H), 6.30(m, 1H), 6.60(s, 1H), 7.08(d, 1H), 7.88(dd,

1H), 8.79(d, 1H), 10.79(brs, 1H).

MS(ES⁺): m/z 533 (M+NH₄).

Compound 12:

2-{2-ethoxyl-5-[2-(pyrrolidinyl-1)ethylamino-N-sulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and 2-(pyrrolidinyl-1)ethylamine.

¹H NMR(CDCl₃) δ: 0.89(t, 3H), 1.59(t, 3H), 1.64(brs, 4H), 1.81(q, 2H), 2.33(brs, 4H), 2.41(s, 3H), 2.52(t, 2H), 3.01(t, 2H), 4.09(t, 2H), 4.33(q, 2H), 6.60(s, 1H), 7.09(d, 1H), 7.89(dd, 1H), 8.84(s, 1H).

MS(ES⁺): m/z 505 (M+NH₄).

Compound 13:

2-[2-ethoxyl-5-(morpholino-4-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

2-[2-ethoxyl-5-(morpholino-4-sulfonyl)phenyl]-5-methyl-3,7-dihydro pyrrolo[2,3-d]pyrimidin-4-one (IG) was prepared from the reaction of 2-amino-3-cyano-4-methylpyridine (IE) and 2-ethoxyl-1-(morpholino-4-sulfonyl) benzoyl chloride (IF) via the corresponding intermediate (ID). The resulting compound was reacted with n-propyl bromide (IH) to give the titled compound.

¹H NMR(CDCl₃) δ: 0.91(t, 3H), 1.64(t, 3H), 1.80(m, 2H), 2.42(s, 3H), 3.03(m, 4H), 3.74(m, 4H), 4.10(t, 2H), 4.39(q, 2H), 6.66(s, 1H), 7.15(d, 1H), 7.80(dd, 1H), 8.77(d, 1H), 10.89(brs, 1H).

MS (ES⁺): m/z 478 (M+NH₄).

Compound 14:

2-[2-ethoxyl-5-(3-(morpholino-4)-n-propylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

5 The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and 3-(morpholino-4)-n-propylamine.

¹H NMR(CDCl₃) δ: 0.92 (t, 3H), 1.59 (t, 3H), 1.72 (m, 2H), 1.84 (q, 2H),
10 2.43 (s, 3H), 2.40-2.50 (m, 6H), 3.11 (t, 2H), 3.72 (m, 4H), 4.11 (t, 2H), 4.32 (q, 2H), 6.60 (s, 1H), 7.13 (t, 1H), 7.88 (d, 1H), 8.79 (s, 1H), 9.95 (brs, 1H).

MS (ES⁺): m/z 535 (M+NH₄).

Compound 15:

15 2-[2-ethoxyl-5-(2-(morpholino-4)-ethylamino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and
20 2-(morpholino-4)ethylamine.

¹H NMR(CDCl₃) δ: 0.90 (t, 3H), 1.58 (t, 3H), 1.80 (m, 2H), 2.32 (m, 4H),
2.44 (s, 3H), 2.74 (t, 2H), 3.03 (t, 2H), 3.58 (m, 4H), 4.06 (t, 2H),
4.30 (q, 2H), 6.62 (s, 1H), 7.08 (d, 1H), 7.87 (d, 1H), 8.79 (d, 1H),
10.82 (brs, 1H).

25 MS (ES⁺): m/z 521 (M+NH₄).

Compound 16:

2-[2-ethoxyl-5-(2,6-dimethylmorpholino-N-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

2-[2-ethoxyl-5-(2,6-dimethylmorpholino-4-sulfonyl)phenyl]-5-methyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one (IG) was prepared from the reaction of 2-amino-3-cyano-4-methylpyrrole (IE) and 2-ethoxyl-5-(2,6-dimethylmorpholino-4-sulfonyl) benzoyl chloride (IF) via the corresponding intermediate ID. The resulting compound was reacted with n-propyl bromide (IH) to give the titled compound.

¹H NMR(CDCl₃) δ: 0.87 (t, 3H), 1.09 (s, 3H), 1.11 (s, 3H), 1.61 (t, 3H), 1.80 (m, 2H), 1.96 (t, 2H), 2.39 (s, 3H), 3.53(s, 1H), 3.58 (s, 1H), 3.70 (m, 2H), 4.08 (t, 2H), 4.33 (q, 2H), 6.60 (s, 1H), 7.10 (d, 1H), 7.77 (dd, 1H), 8.69 (d, 1H), 10.76 (br, 1H).

MS(ES⁺): m/z 506 (M+NH₄).

Compound 17:

2-[2-ethoxyl-5-(benzylpiperidyl-4-aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 2-ethoxyl-5-(1-benzylpiperidyl-4-aminosulfonyl) benzoyl chloride (IF) and 1-n-propyl-2-amino-3-cyano-4-methylpyrrole(IE).

¹H NMR(CDCl₃) δ: 0.82 (t, 3H), 1.50-1.80 (m, 9H), 2.01(m, 2H), 2.40 (s, 3H), 2.76 (m, 2H), 3.14 (m, 1H), (s, 2H), 3.99(t, 2H), 4.23 (q, 2H), 6.58 (s, 1H), 6.94 (d, 1H), 7.21-7.27 (m, 5H), 7.84 (dd, 1H), 8.75 (d, 1H).

MS(ES⁺): m/z 581 (M+NH₄).

Compound 18:

2-[2-ethoxyl-5-(2-(piperidin-1-yl)ethylamine-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) of Preparation 5 and 2-(piperidin-1-yl)ethylamine.

^1H NMR(CDCl_3) δ : 0.89(t, 3H), 1.40(m, 2H), 1.50-1.70(m, 7H), 1.79(m 2H), 2.38(s, 3H), 2.57(brs, 4H), 2.65(t, 2H), 3.07(t, 2H), 4.08(t, 2H), 4.29(q, 2H), 6.61(s, 1H), 7.08(d, 1H), 7.87(d, 1H), 8.77(s, 1H).

10 MS(ES^+): m/z 519 ($\text{M}+\text{NH}_4$).

Compound 19:

2-[2-ethoxyl-5-(4-benzylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 2-ethoxyl-5-(4-benzylpiperazinyl-1-sulfonyl) benzoyl chloride (IF) and 1-n-propyl-2-amino-3-cyano-4-methylpyrrole (IE) via intermediate ID.

^1H NMR(CDCl_3) δ : 0.97(t, 3H), 1.62(t, 3H), 1.80(m, 2H), 2.41(s, 3H), 2.60(m, 4H), 2.63(s, 2H), 3.09(m, 4H), 4.07(t, 2H), 4.35(q, 2H), 6.61(s, 1H), 7.10(d, 1H), 7.21-7.29(m, 5H), 7.78(dd, 1H), 8.78(d, 1H), 10.64(brs, 1H).

20 MS(ES^+): m/z 567 ($\text{M}+\text{NH}_4$).

Compound 20:

2-[2-ethoxyl-5-(4-phenylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

25 1-n-propyl-2-(2-ethoxyl-5-(4-phenylpiperazinyl-1-sulfonyl)benzoyl

amino)-4-methyl-1H-pyrrolo-3-formamide (1A) was prepared from the reaction of 1-n-propyl-2-amino-3-cyano-4-methylpyrrole (IE) and 2-ethoxyl-5-(4-benzylpiperidinyl-1-sulfonyl) benzoyl chloride (IF) via the intermediate ID. A cyclization reaction of IA was effectively carried out to give the titled compound.

^1H NMR (CDCl_3) δ : 0.91 (t, 3H), 1.64 (t, 3H), 1.82 (m, 2H), 2.41 (s, 3H), 3.29 (m, 8H), 4.11 (t, 2H), 4.36 (q, 2H), 6.63 (s, 1H), 6.84-7.05 (m, 2H), 7.14 (d, 1H), 7.20-7.30 (m, 3H), 7.84 (dd, 1H), 8.82 (d, 1H), 10.64 (brs, 1H).

MS (ES^+): m/z 553 ($\text{M}+\text{NH}_4$).

10 Compound 21:

2-[2-ethoxyl-5(piperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) and excess amount of piperazine.

^1H NMR (CDCl_3) δ : 0.91 (t, 3H), 1.60 (t, 3H), 1.77-1.86 (m, 2H), 2.41 (s, 3H), 2.47 (brs, 4H), 2.96 (brs, 4H), 4.08 (t, 2H), 4.29-4.35 (q, 2H), 6.61 (s, 1H), 7.14 (d, 1H), 7.80 (dd, 1H), 8.70 (d, 1H), 10.68 (s, 1H).

20 MS (ES^+): m/z 477 ($\text{M}+\text{NH}_4$).

Compound 22:

2-[2-ethoxyl-5-(4-benzo[1,3]dioxol-5-yl-methylpiperazinyl-1-sulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

25 1-n-propyl-2-(2-ethoxyl-5-(4-benzo[1,3]dioxol-5-yl-methylpiperaziny

1-1-sulfonyl)benzoylamino)-4-methyl-1H-pyrrolo-3- formamide (IA) was prepared from the reaction of 1-n-propyl-2-amino-3-cyano-4-methylpyrrole (IE) and 2-ethoxyl-5-(4-benzo[1,3]dioxol-5-yl-methylpiperazinyl-1-sulfonyl)benzoyl chloride (IF) via the intermediate ID. A cyclization reaction of IA was effectively carried out to give the titled compound.

^1H NMR(CDCl_3) δ : 0.87(t, 3H), 1.63(t, 3H), 1.76(m, 2H), 2.41(s, 3H), 2.51(brs, 4H), 3.08 (brs, 4H), 3.38(s, 2H), 4.08 (t, 2H), 4.34(q, 2H), 5.89(s, 2H), 6.61(s, 1H), 6.62-6.80(m, 3H), 7.11(d, 1H), 7.75(dd, 1H), 8.76(d, 1H), 10.64(s, 1H).

MS(ES^+): m/z 611 ($\text{M}+\text{NH}_4$).

Compound 23:

2-{2-ethoxyl-5-[4-(3-phenyl-n-propan-1-yl)piperidyl-1-sulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and 4-(3-phenyl-n-propan-1-yl) piperidine.

^1H NMR(CDCl_3) δ : 0.90(t, 3H), 1.23-1.31(m, 5H), 1.50-1.85(m, 9H), 2.25(t, 2H), 2.41(s, 3H), 2.52(m, 2H), 3.77 (d, 2H), 4.09(t, 2H), 4.34(q, 2H), 6.61(s, 1H), 7.08-7.29(m, 6H), 7.80(dd, 1H), 8.78(d, 1H).

MS(ES^+): m/z 594 ($\text{M}+\text{NH}_4$).

Compound 24:

2-[2-ethoxyl-5-(n-propylamino-1-sulfonyl)phenyl]-5-methyl-7-n-

propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the reaction of 1-n-propyl-2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA and IC. The resulting compound was reacted with n-propylamine to give the titled compound.

^1H NMR(CDCl_3) δ : 0.94-0.99(m, 6H), 1.52(q, 2H), 1.61(t, 3H), 1.77(m, 2H), 2.40(s, 3H), 2.92(brs, 2H), 4.09(t, 2H), 4.34(q, 2H), 4.82(brs, 1H), 6.64(s, 1H), 7.08(d, 1H), 7.89(dd, 1H), 8.83(d, 1H), 10.90(brs, 1H).

MS(ES^+): m/z 450 ($\text{M}+\text{NH}_4$).

Compound 25:

2-[2-ethoxyl-5-(N,N-di(2-hydroxyethyl)aminosulfonyl)phenyl]-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the reaction of 1-n-propyl-2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA and IC. The resulting compound was reacted with N,N-di(2-hydroxyethyl)amine to give the titled compound.

^1H NMR(CDCl_3) δ : 0.90(t, 3H), 1.58(t, 3H), 1.78(m, 2H), 2.39(s, 3H), 3.32(t, 4H), 3.85(t, 4H), 4.10(t, 2H), 4.35(q, 2H), 6.64(s, 1H), 7.10(d, 1H), 7.85(dd, 1H), 8.79(d, 1H), 10.84(brs, 1H).

MS(ES^+): m/z 496 ($\text{M}+\text{NH}_4$).

Compound 26:

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-methyl]aminosulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the
5 reaction of 1-n-propyl-2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA and IC. The resulting compound was reacted with N-(2-hydroxyethyl)-N-methylamine to give the titled compound.

¹H NMR(CDCl₃) δ: 0.90(t, 3H), 1.62(t, 3H), 1.78(m, 2H), 2.41(s, 3H),
10 2.87(s, 3H), 3.20(t, 2H), 3.77(t, 2H), 4.10(t, 2H), 4.35(q, 2H), 6.65(s, 1H), 7.14(d, 1H), 7.85(dd, 1H), 8.79(d, 1H), 10.89(brs, 1H).

MS(ES⁺): m/z 466 (M+NH₄).

Compound 27:

2-{2-ethoxyl-5-[N-(2-hydroxyethyl-N-ethyl)aminosulfonyl]phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:
15

4-ethoxyl-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the
reaction of 1-n-propyl-2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID,
20 IA and IC. The resulting compound was reacted with N-(2-hydroxyethyl)-N-ethylamine to give the titled compound.

¹H NMR(CDCl₃) δ: 0.90(t, 3H), 1.18(t, 3H), 1.62(t, 3H), 1.79(m, 2H),
2.41(s, 3H), 3.30(m, 4H), 3.75(t, 2H), 4.08(t, 2H), 4.32(q, 2H), 6.61(s, 1H), 7.10(d, 1H), 7.86(d, 1H), 8.81(d, 1H), 10.69(brs, 1H).

25 MS(ES⁺): m/z 480 (M+NH₄).

Compound 28:

2-{2-ethoxyl-5-[N-(2-hydroxyethyl)-N-n-butyl]aminosulfonyl}phenyl}-
5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one

The titled compound was prepared from the reaction of
5 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]
pyrimidin-2-yl)benzenesulfonylchloride (IB) of Preparation 5 and
N-(2-hydroxyethyl)-N-n-butylamine.

^1H NMR(CDCl_3) δ : 0.85-0.93(m, 6H), 1.29-1.40(t, 2H), 1.49-1.65(m, 35),
1.82, 2H), 2.40(s, 3H), 3.15-3.30(m, 4H), 3.78(t, 2H), 4.11(t, 2H),
10 4.38(q, 2H), 6.66(s, 1H), 7.12(d, 1H), 7.88(dd, 1H), 8.82(d, 1H),
10,96(brs, 1H).

MS(ES^+): m/z 508 ($\text{M}+\text{NH}_4$).

Compound 29:

2-{2-ethoxyl-5-(p-ethoxylcarboxylphenylamino)-N-sulfonyl}phenyl}-5-
15 methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

2-[2-ethoxyl-5-(p-ethoxylcarboxylphenylamino-N-sulfonyl)phenyl]-5-
methyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one (IG) was prepared
from the reaction of 2-amino-3-cyano-4-methylpyrrole (IE) and
2-ethoxyl-5-(p-ethoxylcarboxylphenylamino)-N-sulfonyl benzoyl
20 chloride (IF) via the corresponding intermediates ID and IA. The
resulting compound was reacted with n-propyl bromine (IH) to give the
titled compound.

^1H NMR(CDCl_3) δ : 0.82(t, 3H), 1.31(t, 3H), 1.49(t, 3H), 1.74(m, 2H),
2.40(s, 3H), 4.39(t, 2H), 4.16-4.32(m, 4H), 6.60(m, 3H), 6.90(d, 1H),
25 7.25(d, 2H), 7.79-7.90(m, 3H), 8.70(d, 1H), 8.97(s, 1H), 10.83(s, 1H).

MS(ES^+): m/z 566 ($\text{M}+\text{NH}_4$).

Compound 30:

2-{2-ethoxyl-5-(o-benzoylphenylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

The titled compound was prepared from the reaction of
5 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and o-benzoyl phenylamine .

¹H NMR(CDCl₃) δ: 0.91(t, 3H), 1.52(t, 3H), 1.78(m, 2H), 2.41(s, 3H),
4.02(t, 2H), 4.13(m, 2H), 6.63(s, 1H), 6.80(d, 1H), 7.0-7.55(m, 7H),
10 7.72(dd, 1H), 7.80(d, 2H), 8.54(d, 1H), 9.79(s, 1H), 10.62(brs, 1H).

MS(ES⁺): m/z 588 (M+NH₄).

Compound 31:

2-{2-ethoxyl-5-(N2-acethydrazido)-N1-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

15 The titled compound was prepared from the reaction of
4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) of Preparation 5 and acethydrazide.

¹H NMR(CDCl₃) δ: 0.90(t, 3H), 1.56(t, 3H), 1.79(m, 2H), 1.90(s, 3H),
20 2.39(s, 3H), 4.09(t, 2H), 4.27(q, 2H), 6.67(s, 1H), 7.01(d, 1H), 7.52(brs, 1H), 7.9(dd, 1H), 8.74(m, 2H).

MS(ES⁺): m/z 465 (M+NH₄).

Compound 32:

2-{2-ethoxyl-5-(2-dimethylaminoethylamino)-N-sulfonyl}phenyl}-5-methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:
25

The titled compound was prepared from the reaction of 4-ethoxyl-3-(5-methyl-4-oxy-7-n-propyl-4,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)-benzenesulfonylchloride (IB) obtained from previous Preparation 5 and 2-dimethylaminoethylamine.

5 ^1H NMR(CDCl_3) δ : 0.90(t, 3H), 1.61(t, 3H), 1.80(m, 2H), 2.09(m, 6H), 2.36(t, 2H), 2.40(s, 3H), 3.02(t, 2H), 4.10(t, 2H), 4.34(q, 2H), 6.61(s, 1H), 7.11(d, 1H), 7.90(dd, 1H), 8.83(d, 1H).

MS(ES^+): m/z 479 ($\text{M}+\text{NH}_4$).

Compound 33:

10 2-{2-ethoxyl-5-(4-ethylpiperazinyl-1-sulphonyl)phenyl}-5-ethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-ethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the reaction of 2-amino-3-cyano-4-ethylpyrrole (IE) and 2-ethoxyl benzoyl
15 chloride (IF) via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

^1H NMR(CDCl_3) δ : 0.90-0.95(m, 6H), 1.56-1.60(m, 6H), 1.75(m, 2H), 2.49(brs, 4H), 2.53(q, 2H), 3.06(m, 4H), 4.05(q, 2H), 4.34(q, 2H), 6.61(s,
20 1H), 7.10(d, 1H), 7.75(dd, 1H), 8.79(d, 1H), 9.90(br, 1H).

MS(ES^+): m/z 519 ($\text{M}+\text{NH}_4$).

Compound 34:

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulphonyl)phenyl]-5-morpholinomethyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

25 4-ethoxyl-3-(5-morpholinomethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo

[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the reaction of 2-amino-3-cyano-4-morpholinomethylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with
5 1-ethylpiperazine to give the titled compound.

$^1\text{H NMR}(\text{CDCl}_3)$ δ : 0.90-1.00 (m, 6H), 1.61 (t, 3H), 1.74 (m, 2H), 2.32-2.56 (m, 8H), 3.06 (m, 4H), 3.67 (m, 4H), 3.94 (s, 2H), 4.07 (t, 2H), 4.34 (q, 2H), 6.55 (s, 1H), 7.15 (d, 1H), 7.78 (dd, 1H), 8.75 (d, 1H).

MS(ES^+): m/z 576 ($\text{M}+\text{NH}_4$).

10 Compound 35:

2-[2-ethoxyl-5-(4-ethylpiperazinyl-1-sulphonyl)phenyl]-5-(pyrimidinyl-2)methyl-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:

4-ethoxyl-3-(5-morpholinomethyl-4-oxy-7-n-propyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from
15 the reaction of 2-amino-3-cyano-4-(pyrimidinyl-2)methylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

$^1\text{H NMR}(\text{CDCl}_3)$ δ : 0.90-1.00 (m, 6H), 1.60 (m, 6H), 1.75 (m, 2H), 2.37 (q, 2H),
20 2.53 (m, 4H), 3.06 (m, 4H), 4.01 (s, 2H), 4.07 (t, 2H), 4.35 (q, 2H), 6.69 (s, 1H), 7.11-7.15 (m, 2H), 7.77 (dd, 1H), 8.68-8.71 (m, 3H).

MS(ES^+): m/z 569 ($\text{M}+\text{NH}_4$).

Compound 36:

2-{2-ethoxyl-5-(4-ethylpiperazinyl-1-sulphonyl)phenyl}-5-methyl-7-allyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one:
25

4-ethoxyl-3-(5-methyl-4-oxy-7-allyl-3,7-dihydropyrrolo[2,3-d]pyrimidin-2-yl)benzenesulfonylchloride (IB) was prepared from the reaction of 2-amino-3-cyano-4-methylpyrrole (IE) and 2-ethoxyl benzoyl chloride (IF) via the corresponding intermediates ID, IA, IG (IH) and IC. The resulting compound was reacted with 1-ethylpiperazine to give the titled compound.

^1H NMR(CDCl_3) δ : 1.00(t, 3H), 1.65(t, 3H), 2.37(q, 2H), 2.41(s, 3H), 2.50(m, 4H), 3.09(br, 4H), 4.31(q, 2H), 4.95(m, 2H), 4.96(dd, 1H), 5.01(dd, 1H), 5.68(m, 1H), 6.61(s, 1H), 7.11(d, 1H), 7.82(dd, 1H), 8.71(d, 1H), 9.77(br, 1H).

MS(ES^+): m/z 503 ($\text{M}+\text{NH}_4$).

Example 2

Experiments of penis erection

In order to demonstrate efficacy of the compounds of formula I in treatment of functional impotence, the penis erection experiment was established by using the male rabbit as experimental model cross reference to Bischoff's method. (Bischoff E.; Schneider K, *International Journal of Impotence Research*, 2001, 13, 230 235)

The hydrochloride of 3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one derivatives was dissolved into water, and injected to conscious-rabbits intravenously (i.v.) (0.1mg-3mg/kg). Erection was accessed by comparing the length of penis before and at 0, 30, 60, 90 and 120 minutes after intravenous administration of the above-mentioned agents. The results of typical compounds in this erectile experiment were listed in table 1.

Table 1. penis erection efficacy of 3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one derivatives

Compounds	Length of rabbit penis (mm)				
	0 min	30 min	60 min	90 min	120 min
Compound1-HCl	8.3	5.7	5.3	2.0	1.3
Compound1-2HCl	1.3	3.0	2.0	1.0	1.0
Compound6-2HCl	0.0	5.3	2.7	1.7	0.0
Compound8-2HCl	2.7	1.7	1.3	0.7	0.0
Compound19-2HCl	1.7	4.0	4.3	4.0	0.0

*The data above were the average value of repeated experimental measurements of 3 rabbits.

The results showed that the above 3,7-dihydropyrrolo[2,3-d]pyrimidin-4-one derivatives had the good potential in treatment of functional erectile dysfunction with the characteristics of quick onset and prolonged effectiveness, especially the compound1-1HCl.

Example 3

Experiments of the effect of single administration on the coitus function of male rats

After purchasing, the male rat and the estrogenized female rat were kept in the same cage for 2 days, and the male rat would acquire the sexual experiences. Then, the female rat was taken out of the cage, and the experiments began after the male rat stayed alone in a cage for 5 days. Each tested compound was orally administered in a dosage of 24mg/kg, the positive group was administered with Sildenafil citrate (Viagra) in a dosage of 24mg/kg, and the control group was administered with saline

at the equal volume (0.1ml/10g) . Fifty minutes after administration, the tested male rats were placed within a observation container (both the diameter and the height of the container were 24cm), and the male rats adapted the new environment for 5 minutes. After that, two
5 estrogenized female rats were placed into the container. The sexual behavior of the male rats in 20 minutes was observed under the non-interfering condition using Panasonic WVCP410/G monitor, the latent period of straddle, times of straddle, as well as the latent period of the coitus and times of the coitus. All the experiments were carried
10 out at 21-24°C, and completed before 11:00 AM. The comparative results among the compounds of the invention, the blank and the positive control drug Sildenafil citrate were listed in Table 2.

The experimental results showed that all the indexes of sexual function of the rats were obviously enhanced after they were administered with
15 compound 1-HCl, namely, the latent period of straddle and coitus were remarkably shortened, and the times of the coitus was obviously increased. And the effect of the compound 1-HCl was stronger than that of the Sildenafil citrate group, especially, the times of the coitus was much more than that of the positive control group($p < 0.05$). All the indexes
20 of sexual function of the rats were also obviously enhanced after they were orally administered with Compound 6-HCl, in particular, the latent period of coitus was remarkably shorter than that of the positive control group. Therefore, the compound of the general formula I of the present invention can obviously enhance the sexual function of rats, and its
25 effect was stronger than that of Sildenafil citrate.

Table 2. Effect of 3,7-dipyrrolo[2,3-d]pyrimidin-4-one derivatives on the indexes of sexual function of the male rats (average values \pm standard deviations)

Groups	Number of rats	the latent period of straddle (min)	times of straddle	the latent period of the coitus (min)	times of the coitus
saline	17	8.9 ± 3.69	7.1 ± 3.62	16.2 ± 4.02	3.4 ± 4.64
Compound1-HCl	9	$2.6 \pm 1.13^{**}$	7.6 ± 2.70	$7.1 \pm 5.62^{**}$	$13.1 \pm 8.77^{**\#}$
Compound6-HCl	8	6.4 ± 3.07	8.0 ± 3.93	$8.8 \pm 5.85^{*}$	9.0 ± 8.50
Sildenafil citrate	19	$5.0 \pm 4.74^{*}$	12.7 ± 7.18	$8.2 \pm 4.02^{**}$	$8.2 \pm 5.25^{*}$

Compared with the group of saline, $*P < 0.05$, $**P < 0.01$; Compared with the group of Sildenafil citrate, $\#P < 0.05$.

Example 4

5 The bioactivity inhibiting phosphodiesterase of compounds of formula I

The bioactivity inhibiting phosphodiesterase 5 (PDE5) of compounds of formula I was measured cross the reference of methods (Hidaka H, et al Biochim. Biochim. Biophys. Acta, 1976, 429, 485; Kim D-K, et al., Bioorg. Med. Chem. 2001, 9, 3013). The inhibiting activity of PDE5 was determined using SPA technique and the method of chemical fluorescence. The method was as follows: firstly, the reaction time curve and the enzyme concentration curve of PDE5 reaction system were determined by Microplate Scintillation & Luminescence Counter (TopCount Counter), by
15 which the optimal reaction condition was determined. Under the optimized

conditions, the inhibition experiments on PDE5 were carried out. The results showed that compound I had stronger inhibiting rate on phosphodiesterase 5 than that of sildenafil. For example, when the concentration of compound 1-HCl was 10^{-8} mol/L, the inhibiting rate of compound 1-HCl on PDE5 was 65.62%, but the inhibiting rate of sildenafil on PDE5 was 31.67%.